

***In vivo* catalyzed new-to-nature reactions**

Johannes G. Rebelein and Thomas R. Ward*

Address

Department of Chemistry, University of Basel, BPR 1096, Mattenstrasse 24a, CH-4058 Basel, Switzerland

*corresponding author: Ward, Thomas R. thomas.ward@unibas.ch

Abstract

Bioorthogonal chemistry largely relies on the use of abiotic metals to catalyze new-to-nature reactions in living systems. Over the past decade, metal complexes and metal-encapsulated systems such as nanoparticles have been developed to unravel the reactivity of transition metals, including ruthenium, palladium, iridium, copper, iron, and gold in biological systems. Thanks to these remarkable achievements, abiotic catalysts are able to fluorescently label cells, uncage or form cytotoxic drugs and activate enzymes *in cellulo/vivo*. Recently, strategies for the delivery of such catalysts to specific cell types, cell compartments or proteins were established. These studies reveal the enormous potential of this emerging field and its application in both medicinal chemistry and in synthetic biology.

Introduction

Over the past decade, abiotic transition metal-catalyzed reactions have successfully been introduced into living cells and organisms. With *in vivo* applications in mind, several challenges were identified and addressed: i) overcome the inherent cytotoxicity of the abiotic catalyst, ii) ensure the efficient metal uptake by the cell and iii) circumvent the deactivation of the catalyst by thiols, proteins and other cell components [1]. To address these issues, several strategies were developed: i) metals were incorporated into nanoparticles or embedded in microspheres and resins (see sections: “Palladium nanoparticles” and “Copper and gold nanoparticles”; the catalyzed reactions are summarized in Table 1) or ii) used as homogeneous metal complexes (see sections: “Homogenous ruthenium catalysts”, “Homogenous palladium catalysts” and “Homogenous iron, iridium and gold catalysts”; the catalyzed reactions and complexes are collected in Table 2). Most reactions developed to date are based on either a cleavage, or a cross-coupling reaction. As a result, a variety of *in vivo* functions were established that are schematically presented in Figure 1 including: i) fluorescent labeling of cells, cell compartments and proteins, ii) synthesis of cytotoxic agents and iii) enzyme rescue. Several excellent reviews cover bioorthogonal reactions [2-7] or bioorthogonal protein labeling and protein chemistry [8-12]. This review focuses on bioorthogonal transition metal-catalyzed reactions *in cellulo* (inside single cells in a cell culture) and *in vivo* (inside living organisms), highlighting recent developments towards therapeutic applications.




Figure 1. Strategies for the introduction of metal-catalyzed bioorthogonal reactions *in cellulo/vivo*. Biocompatible transition metal catalysts including metal-nanoparticles (NPs), NPs entrapped in microspheres or resins, and metal complexes are taken-up by cells or introduced into organisms. These metal-platforms convert *in cellulo/vivo* various substrates through coupling or uncaging reactions into fluorophores, drugs and active enzymes. Recent progress allows to direct metal complexes to specific cell-types or cell compartments such as organelles.

Palladium nanoparticles.

The use of metal nanoparticles (NPs) for *in cellulo* catalysis was first demonstrated by the Bradley group [13]. They exploited the biocompatibility of polystyrene microspheres [14] by entrapping Pd-NPs, creating fluorescently-labelled Pd⁰-microspheres [13,15]. These Pd⁰-catalysts cleave allyloxycarbonyl (alloc)-groups, leading to the uncaging of fluorescent rhodamine 110 (R110) within the cytoplasm of HeLa cells (Table 1, entry 1) [13]. Moreover, these Pd⁰-loaded microspheres also catalyze the Suzuki-Miyaura cross-coupling inside HeLa cells, leading to the accumulation of a rhodamine-fluorophore within mitochondria (Table 1, entry 9) [13].

These findings suggest that such Pd-NPs may find applications, both, in labeling studies as well as in medical applications whereby one can specifically target an organelle. With this goal in mind, Broceta and Bradley *et al.* implemented a Pd-NP catalyzed drug release. Pd⁰-NPs were linked to polyethylene glycol (PEG)-polystyrene resins forming micrometer-sized beads (~150 µm) with a remarkable biocompatibility [16]. A single resin bead was implanted into the yolk sac of zebrafish embryos. The embryos developed normally without any signs of toxicity. To prove the *in vivo* activity of the Pd⁰-resin, the uncaging of a propargyloxycarbonyl (poc)-protected R110 was assayed within the yolk sac (Table 1, entry 3). This Pd⁰ resin was also able to extracellularly uncage 5-fluoro-1-propargyl (pro)-uracil (pro-5FU). This led to the release of the cytotoxic drug 5-fluorouracil (5FU) in a cancer cell culture of either HCT116 or BxPC-3 cells (Table 1, entry 6) [16].

This therapeutic strategy was refined by combining a dual drug synthesis strategy with tumor specificity [17]. To achieve cell specificity, fluorescent Pd-microspheres were decorated with the cancer targeting motif cyclic-RGD (cRGD). cRGD is an antagonist of the α_vβ₃ receptor, an angiogenesis factor overexpressed in many tumors and tumor vasculature.

These tumor-targeting microspheres were used for the activation of two prodrugs inside U87-MG glioblastoma cells. Besides uncaging pro-5FU, the cytotoxic agent PP-121 was produced via a Suzuki-Miyaura cross coupling reaction leading to increased cell death, compared to the synthesis of either 5FU or PP-121 alone (Table 1, entry 6 and 10) [17]. This approach illustrates the potential of metal-NPs for targeted drug delivery by converting innocuous substrates into active drugs inside tumor cells.

Subsequently, Miller *et al.* explored a Pd-NP based drug synthesis strategy in mice [18]. For this purpose, the palladium complex bis[tri(2-furyl)phosphine]palladium(II)-dichloride ($\text{PdCl}_2(\text{TFP})_2$) was encapsulated into poly(lactic-co-glycolic acid)-polyethyleneglycol (PLGA-PEG)-NPs, a material which is in clinical trials for drug administration [19]. The authors thoroughly studied the effect of the nano-encapsulation, observing an improved solubility, bioavailability, biocompatibility and a low cytotoxicity (IC_{50} of 70-130 μM). Furthermore, Pd-NPs were enriched in tumor tissue of cancer mouse models compared to other organs such as brain, skin, lung, bone marrow and heart. Interestingly, the highest catalytic activity of Pd-NPs for the release of either fluorescent R110 or the chemotherapeutic drug doxorubicin was observed in tumor tissue (Table 1, entry 1 and 2). Therefore, combining nano-encapsulated Pd-NPs and caged doxorubicin helped to minimize systemic exposure and toxicity, thus increasing the maximum-tolerated dose by 300 %, resulting in the halt of tumor growth and extending the survival of tumor-bearing mice [18].

Copper and gold nanoparticles.

The range of metal-NPs was recently broadened to include Cu [20,21] and Au [22,23], thus expanding the *in vivo* catalytic repertoire. Although Cu^I-catalyzed azide-alkyne cycloaddition (CuAAC) has been known since 2002 [24,25], it was hardly used *in vivo* due to its marked cytotoxicity [26]. The incorporation of Cu into aspartate-containing polyolefins [20] or entrapping Cu in TentaGel resin [21], contributed to overcoming these biocompatibility issues. On the one hand, Cu-based NPs convert coumarin-derivatives via CuAAC into fluorophores inside NCI-H460 and MDA-MB-231 cancer cells (Table 1, entry 12) [20]. On the other hand, Cu-particles produce *in cellulo* cytotoxic agents, decreasing the cell viability of *Escherichia coli* (Table 1, entry 14) [20]. Furthermore, the implantation of the Cu-resin into the yolk sac of zebrafish embryos did not exhibit any toxicity but catalyzed the release of a coumarin-fluorophore (Table 1, entry 13) [21].

Besides Cu and Pd, gold was immobilized on a solid support, a polyethylene glycol (PEG)-grafted low-cross-linked polystyrene matrix [22]. This Au-resin was employed for the local release of R110 in the brain of zebrafish (Table 1, entry 3) as well as for the activation of structurally diverse prodrugs inside A549 cancer cells (Table 1, entry 5, 7 and 8). This demonstrates the potential application of Au-resin for *in vivo* drug-release [22]. Furthermore, Tonga *et al.* created controllable, Au-NPs by blocking the access to the Au-encapsulated, catalytic Ru- or Pd-complexes [23]. This 'gate-keeping' was achieved through steric hindrance by supramolecular binding of cucurbit[7]uril onto the surface. The NPs were activated through the addition of a competitor, 1-adamantylamine to unmask alloc-R110 (by Ru- and Pd-loaded NPs) and the prodrug pro-5FU (by Pd-loaded NPs) inside live HeLa cells (Table 1, entry 1 and 6) [23]. These gated NPs reduce the interference with cell components, protect the metal from poisoning and increase the biocompatibility. Moreover, the introduction of a switch into these NPs allows a more controlled drug treatment and suggests their use for restoring malfunctioning pathways in chronic disease.

Table 1: Reactions catalyzed by metal loaded nanoparticles















Entry	Reaction	Catalyst	Organism	Ref.
Allyloxycarbonyl cleavage				
1	 <p style="text-align: center;">Rhodamine 110</p>	<p>Entrapped Pd-NPs [13,15]</p> <p>Free Pd-NPs [18]</p> <p>Ru/Pd-complex entrapped in Au-NPs [23]</p>	<p>HeLa cells</p> <p>HT1080, ID8, A2780CP cells; mouse model; HeLa cells</p>	
2	 <p style="text-align: center;">Doxorubicin</p>	Free Pd-NPs	Mouse model	[18]
Propargyloxycarbonyl cleavage				
3	 <p style="text-align: center;">Rhodamine 110</p>	<p>Entrapped Pd-NPs [16]</p> <p>Entrapped Au-NPs [22]</p>	<p>Zebrafish</p> <p>Zebrafish</p>	
4	 <p style="text-align: center;">Cresyl violet</p>	Entrapped Pd-NPs	U87-MG cells	[17]
5	 <p style="text-align: center;">Doxorubicin</p>	Entrapped Au-NPs	A549 cells	[22]
Propargyl cleavage				
6	 <p style="text-align: center;">5FU</p>	<p>Entrapped Pd-NPs [16]</p> <p>Pd-complex entrapped in Au-NPs [23]</p>	<p>HCT116, BxPC-3 cells; U87-MG cells; HeLa cells</p>	
7	 <p style="text-align: center;">FUDR</p>	Entrapped Au-NPs	A549 cells	[22]

Table 1 (continued)

Entry	Reaction	Catalyst	Organsim	Ref.
Propargyloxyaryl cleavage				
8	 <p>Vorinostat</p>	Entrapped Au-NPs	A549 cells	[22]
Suzuki-Miyaura cross-coupling				
9		Entrapped Pd-NPs	HeLa cells	[13,15]
10	 <p>PP-127</p>	Entrapped Pd-NPs	U87-MG cells	[17]
Heck Coupling				
11		Free Pd-NPs	HT80 cells	[18]
Azide-alkyne cycloaddition				
12		Free Cu-NPs	NCI-H460, MDA-MB-231 cells;	[20]
13		Entrapped Cu-NPs	Zebrafish	[21]
14	 <p>Bisamidine</p>	Free Cu-NPs	<i>E. coli</i>	[20]

Homogenous ruthenium catalysts

Streu's and Meggers' seminal work on *in cellulo* catalyzed Ru-reactions laid the foundation of bioorthogonal organometallic chemistry [27]. In 2006, they suggested that the complex [Cp*Ru(cod)Cl] **Ru1** (Cp*=pentamethylcyclopentadienyl, cod=1,5-cyclooctadiene, Table 2, entry 1) catalyzes the uncaging of alloc-protected R110 inside HeLa cells, while not affecting the cell viability [27]. In 2014 and 2017, they reported on significantly more active Ru-pianostool complexes **Ru2-Ru6** (Table 2, entry 1 and 4) catalyzing alloc-unmasking [28,29]. Moreover, alloc-protecting groups were cleaved off from DNA-binding agents such as 4',6-diamidino-2-phenylindole (DAPI) and ethidium bromide inside chicken embryo fibroblasts (CEF) and the kidney epithelial cell line Vero (Table 2, entry 2 and 3) [30].

Subsequently, the quinolone based Ru-complex was improved by reducing the π -backbonding of the bidentate ligand (**Ru2-Ru4**) and improving the *in vivo* R110 release by 130-fold, comparing **Ru4** to **Ru1** [28]. Furthermore, **Ru4** unmasked efficiently alloc-doxorubicin, triggering the apoptosis of HeLa cells [28]. Very recently, the activity of the Ru-complex was further enhanced tenfold by replacing the quinoline of **Ru4** with an 8-hydroxyquinolate motif (**Ru5** and **Ru6**) [29]. **Ru6** decages almost quantitatively alloc-doxorubicin decreasing the IC₅₀ concentration of alloc-doxorubicin from 15 μ M to 2 μ M in the presence of 1 μ M **Ru4** or **Ru6**, respectively [29]. ICP-MS studies reported by Wender *et al.* however suggest that the Ru-complex **Ru3** does not penetrate into the cytoplasm of 4T1 cells, indicating that the reaction occurs extracellularly [31]. To overcome this shortcoming, Mascareñas *et al.* appended a cation triphenylphosphonium (TPP) moiety to **Ru2** to afford **Ru7** (Table 2, entry 1) [32]. Such lipophilic cations are taken up by mitochondria [33,34]. After treading HeLa cells with **Ru7** and caged-R110, fluorescence was measured within mitochondria. This suggests that **Ru7** is present in the mitochondria, thanks to the TPP targeting moiety [32]. To improve the traceability of **Ru7**, a pyrene derivative was added to afford **Ru8** (Table 2, entry 1), enabling fluorescent monitoring of the complex itself. As anticipated, both the catalyst **Ru8** and R110 were localized inside the mitochondria. The authors specifically depolarized mitochondria by deprotecting the proton ionophore 2,4-dinitrophenol (Table 2, entry 5) [35,36], demonstrating the potential of organelle-specific complexes for disturbing and modifying biological systems [32].

A different approach to target cell compartments and proteins was pursued by the Winssinger lab [37-39]. A Ru-complex (**Ru10**, Table 2, entry 6) was linked to a peptide nucleic acid (PNA)-probe which hybridizes with tissue specific miRNAs [39]. The **Ru10**-PNA and a fluorogenic rhodamine-PNA were injected into a one-cell zebrafish embryo. Using PNA-recognition sequences on both the catalyst and the azide-bearing substrate ensures the photocatalytic reduction of the azide upon docking of the substrate and catalyst to the target miRNA. This yields fluorescence in the specific tissues of live zebrafish. The same Ru-complex conjugated to an epidermal growth factor receptor (EGFR)-inhibitor, **Ru11**, and an estrogen agonist (raloxifene), **Ru12**, (Table 2, entry 7) targeted the receptor proteins and provided a spatial and temporal unmasking of the fluorescent dye in live HEK293T and MCF-7 cells [38].

By targeting a biotinylated Ru-cofactor **Ru13** to streptavidin (Sav) in *E. coli*, The Ward group could evolve an artificial metalloenzyme *in cellulo* (Table 2, entry 8) [40]. Fusing the signal peptide of the outer membrane protein A (OmpA) to Sav lead to the efficient secretion of the overexpressed Sav into the periplasm of *E. coli*. Addition of **Ru13** to these cells allowed the assembly of an artificial metalloenzyme **Ru13**-Sav within the periplasm of *E. coli*. The resulting strain catalyzed the ring closing metathesis (RCM) of an umbelliferone precursor in

the periplasm of *E. coli*. The catalytic performance of **Ru13**-Sav was improved by directed evolution relying on the fluorescence of the formed umbelliferone. The evolved strain was shown to catalyze the RCM of typical diolefin substrates, leading to 660 TONs, outperforming the standard homogeneous RCM catalysts [40].

Homogeneous palladium catalysts

To the best of our knowledge, the first homogeneous Pd-catalyzed reaction *in cellulo* was reported by Lin *et al.* [41]. A copper-free Sonogashira cross-coupling reaction was used to selectively label homopropargylglycine *E. coli* cells. Fluorescein iodide was coupled to the metabolically incorporated homopropargylglycine, by a newly discovered Pd-complex **Pd1** (Table 2, entry 9) [41].

Subsequently, unnatural amino acids (UAAs) were genetically encoded by amber stop codon suppression [42,43] and *p*-iodophenylalanine was incorporated on the external loops of the bacterial pore protein OmpC [44]. The palladium complex **Pd1** coupled a fluorescent boronic acid to the *p*-iodophenylalanine of the OmpC-mutants via Suzuki-Miyaura cross-coupling on the surface of live *E. coli*s (Table 2, entry 12) [44]. Furthermore, proteins carrying an encoded alkyne were cross-coupled via a Sonogashira reaction with a fluorophore bearing a iodophenyl group (Iph-FL-525) using [Pd(NO₃)₂] inside *E. coli* and *Shigella* (Table 2, entry 10) [45].

The Pd-catalyzed fluorescent labeling of UAAs was also expanded to mammalian cells [46]. N^ε-butynyloxycarbonyllysine encoded in the EGFR-protein was cross-coupled to biotin-phenyl iodide using catalyst **Pd1** (Table 2, entry 11). The biotinylated EGFR was monitored by fluorescence through addition of Sav conjugated to Alexa Fluor® 568 [46].

Besides labeling specific proteins *in vivo*, Chen and coworkers were the first to employ Pd-complexes for a gain-of-function study, using a deprotection-strategy [47]. The phosphothreonine lyase OspF was blocked by masking the catalytic Lys134 with a poc-group. The catalyst [Pd(η^3 -allyl)Cl]₂ **Pd3** (Table 2, entry 14) cleaved the poc-OspF and allowed the *in vivo* monitoring its substrate, the Erk kinase, in HeLa cells [47]. This gain-of-function approach was extended by incorporating the UAA allenyl-tyrosine that can be uncaged thanks to homogeneous Pd-catalysts **Pd3-Pd6** (Table 2, entry 15) [48]. The following enzymes were rescued by allenyl-removal through **Pd3-Pd6** in HEK293T cells: *Taq* DNA polymerase, Src kinase and the anthrax lethal factor endopeptidase [48].

Recently, cell-penetrating peptides were linked to a Pd-complex **Pd7** (Table 2, entry 13), improving its internalization (Table 2, entry 13) [49]. The uncaging of poc-R110 in prostate adenocarcinoma (PC-3) cells was demonstrated [49]. The peptide-based platform facilitates the delivery of metal-complexes to specific cell-types, thus opening the possibility of targeted drug delivery [50,51].

Homogenous iron, iridium and gold catalysts

Besides Ru- and Pd-complexes, other metals have been employed for intracellular labeling: An iron(III) meso-tetraarylporphyrin **Fe1** (Table 2, entry 16) was shown to catalyze azide reduction leading to the release of R110 in HeLa cells [52]. The Ir-complex **Ir1** (Table 2, entry 17) reduces an aldehyde leading to the release of a fluorescent Bodipy-OH inside NIH-3T3 cells [53]. A more sophisticated Au-catalyzed approach was developed by Tanaka *et al.* [54]. They relied on an Au-complex linked to coumarin, **Au1** (Table 2, entry 18 and 19) which has marked affinity for albumin. These albumins were coated with glycans through the “RIKEN click” reaction, influencing the trafficking pathways and resulting in an accumulation either

on the liver or the intestine by using (2-6)-disialoglycoalbumin or galactosylglycoalbumin, respectively. These organ-targeted Au-complexes catalyzed an amide bond formation between a propargyl-ester of fluorescent probes (TAMRA-O-pro, Cy7.5-O-pro) and surface-exposed amines of nearby proteins leading to an organ-specific labeling inside living mice [54].

Conclusions

Since 2014, the field of metal-catalyzed bioorthogonal reactions has made significant progress. Initially, it was demonstrated that metal-NPs and metal complexes can uncage or synthesize drugs *in cellulo/vivo*. Subsequently, the specificity of these metal-based platforms was considerably improved. Now, it is possible to target specific organs, cells, cell compartments and certain proteins. Thanks to these efforts, one can envision the possibility of targeting and killing cells as well as gaining abiotic cellular functions.

The exploitation of novel targeting and packaging strategies should further contribute to decreasing the toxicity of the metal catalysts and increasing their specificity. This extends the safety window of metal catalysts *in vivo*, hopefully resulting in the development of new therapies. Beyond clinical applications, one can envisage to design bioorthogonal metabolic pathways *in cellulo/vivo* for the production of high-value chemicals.

Table 2: Reactions catalyzed by metal- complexes in a biological environment









Entry	Reaction	Catalyst	Organism	Ref.
1	 <p>Rhodamine 110</p>	Ruthenium	HeLa cells	[27]
		 <p>Ru1</p>	HeLa cells	[28]
		 <p>Ru2 -H Ru3 -OMe Ru4 -NMe₂</p>	HeLa cells and mitochondria of HeLa cells	[32]
		 <p>Ru7 Ru8 Ru9</p>		
2	 <p>DAPI</p>	 <p>Ru1</p>	CEF, Vero cells	[30]
3	 <p>Ethidium bromide</p>	 <p>Ru1</p>	CEF cells	[30]

Table 2 (continued)













Entry	Reaction	Catalyst	Organism	Ref.
Allyloxycarbonyl cleavage				
4	 Doxorubicin	 Ru2 -H Ru3 -OMe Ru4 -NMe ₂ + PF ₆ ⁻	HeLa cells	[28]
		 Ru5 -H Ru6 -CO ₂ Me + PF ₆ ⁻	HeLa cells	[29]
Allyl cleavage				
5	 2,4-dinitrophenol	 Ru8  Ru9  + PF ₆ ⁻	HeLa cells and mitochondria of HeLa cells	[32]
Photocatalyzed azide reduction				
7		 Ru10	Zebrafish	[39]
6		 Ru11  Ru12	HEK293T, MCF-7 cells	[38]

Table 2 (continued)










Entry	Reaction	Catalyst	Organism	Ref.
Ring closing metathesis				
8	 <p>Umbelliferone</p>	 <p>Ru13</p>	Periplasm of <i>E. coli</i>	[40]
Copper-free Sonogashira cross-coupling				
9	 <p>Fluorescein</p>	 <p>Pd1</p>	<i>E. coli</i>	[41]
10	 <p>Biotin</p>	 <p>Pd1</p>	HEK293	[46]
11	 <p>Fluor 525</p>	$\text{Pd}(\text{NO}_3)_2$ (Pd2)	<i>E. coli</i> and <i>Shigella</i>	[45]
Suzuki-Miyaura cross-coupling				
12		 <p>Pd1</p>	<i>E. coli</i>	[40,44]

Table 2 (continued)












Entry	Reaction	Catalyst	Organism	Ref.
Propargyloxycarbonyl cleavage		Palladium		
13	 <p style="text-align: center;">Rhodamine 110</p>	$[\text{Pd}(\eta^3\text{-allyl})\text{Cl}]_2$ (Pd3) $\text{Pd}(\text{dba})_2$ (Pd4)	HEK293T, NIH-3T3, A549, HeLa, Caco-2, CHO cells PC-3 cells	[47] [41] [49]
14	 <p style="text-align: center;">Lysine</p>	$[\text{Pd}(\eta^3\text{-allyl})\text{Cl}]_2$ (Pd3) $\text{Pd}(\text{dba})_2$ (Pd4)	HeLa cells	[47]
Allenyl cleavage		Palladium		
15	 <p style="text-align: center;">Tyrosine</p>	$[\text{Pd}(\eta^3\text{-allyl})\text{Cl}]_2$ (Pd3) $\text{Pd}(\text{dba})_2$ (Pd4) $\text{Pd}(\text{TAPAd})_2$ (Pd5) $\text{Pd}(\text{TPPTS})_4$ (Pd6)	HEK293T cells	[48]
Bisazide reduction		Iron		
16	 <p style="text-align: center;">Rhodamine 110</p>	 <p style="text-align: center;">Fe1</p>	HeLa cells	[40,52]
Transfer Hydrogenation		Iridium		
17	 <p style="text-align: center;">Bodipy-OH</p>	 <p style="text-align: center;">Ir1</p>	NIH-3T3 cells	[41,53]

Table 2 (continued)

Entry	Reaction	Catalyst	Organism	Ref.
	Amide bond formation	Gold		
18	 <p>Cy7.5</p>	 <p>Au1</p>	Mouse liver and intestine	[54]
19	 <p>TAMRA</p>	 <p>Au1</p>	Mouse liver and intestine	[54]

Acknowledgments: TRW thanks the ERC (The DrEAM) and the NCCR Molecular Systems engineering for generous support of his work in this field. JGR thanks EMBO for a Long-Term fellowship (EMBO ALTF 194-2017).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Wilson YM, Durrenberger M, Nogueira ES, Ward TR: **Neutralizing the detrimental effect of glutathione on precious metal catalysts.** *J Am Chem Soc* 2014, **136**:8928-8932.
2. Sletten EM, Bertozzi CR: **Bioorthogonal chemistry: fishing for selectivity in a sea of functionality.** *Angew Chem Int Ed Engl* 2009, **48**:6974-6998.
3. Sasmal PK, Streu CN, Meggers E: **Metal complex catalysis in living biological systems.** *Chem Commun (Camb)* 2013, **49**:1581-1587.
4. Patterson DM, Nazarova LA, Prescher JA: **Finding the right (bioorthogonal) chemistry.** *ACS Chem Biol* 2014, **9**:592-605.
5. Chankeshwara SV, Indrigo E, Bradley M: **Palladium-mediated chemistry in living cells.** *Curr Opin Chem Biol* 2014, **21**:128-135.
6. Volker T, Meggers E: **Transition-metal-mediated uncaging in living human cells-an emerging alternative to photolabile protecting groups.** *Curr Opin Chem Biol* 2015, **25**:48-54.
7. Li J, Chen PR: **Development and application of bond cleavage reactions in bioorthogonal chemistry.** *Nat Chem Biol* 2016, **12**:129-137.
8. Spicer CD, Davis BG: **Selective chemical protein modification.** *Nat Commun* 2014, **5**:4740.
9. Yang M, Li J, Chen PR: **Transition metal-mediated bioorthogonal protein chemistry in living cells.** *Chem Soc Rev* 2014, **43**:6511-6526.
10. Gong Y, Pan L: **Recent advances in bioorthogonal reactions for site-specific protein labeling and engineering.** *Tetrahedron Lett* 2015, **56**:2123-2132.
11. Chen X, Wu YW: **Selective chemical labeling of proteins.** *Org Biomol Chem* 2016, **14**:5417-5439.
12. Schwizer F, Okamoto Y, Heinisch T, Gu Y, Pellizzoni MM, Lebrun V, Reuter R, Kohler V, Lewis JC, Ward TR: **Artificial Metalloenzymes: Reaction Scope and Optimization Strategies.** *Chem Rev* 2017, doi: [10.1021/acs.chemrev.7b00014](https://doi.org/10.1021/acs.chemrev.7b00014).
- This is a comprehensive review about artificial metalloenzymes (ArMs). Used scaffold proteins, anchoring strategies and metallocofactors are discussed. Mutagenesis approaches for the evolution of ArMs are reviewed. The ArMs are organized by reaction class.
13. Yusop RM, Unciti-Broceta A, Johansson EM, Sanchez-Martin RM, Bradley M: **Palladium-mediated intracellular chemistry.** *Nat Chem* 2011, **3**:239-243.
14. Alexander LM, Pernagallo S, Livigni A, Sanchez-Martin RM, Brickman JM, Bradley M: **Investigation of microsphere-mediated cellular delivery by chemical, microscopic and gene expression analysis.** *Mol Biosyst* 2010, **6**:399-409.

15. Unciti-Broceta A, Johansson EM, Yusop RM, Sanchez-Martin RM, Bradley M: **Synthesis of polystyrene microspheres and functionalization with Pd(0) nanoparticles to perform bioorthogonal organometallic chemistry in living cells.** *Nat Protoc* 2012, **7**:1207-1218.
16. Weiss JT, Dawson JC, Macleod KG, Rybski W, Fraser C, Torres-Sanchez C, Patton EE, Bradley M, Carragher NO, Unciti-Broceta A: **Extracellular palladium-catalysed dealkylation of 5-fluoro-1-propargyl-uracil as a bioorthogonally activated prodrug approach.** *Nat Commun* 2014, **5**:3277.
17. Clavadetscher J, Indrigo E, Chankeshwara SV, Lilienkamp A, Bradley M: **In-Cell Dual Drug Synthesis by Cancer-Targeting Palladium Catalysts.** *Angew Chem Int Ed Engl* 2017, **56**:6864-6868.

••

Pd-microspheres were coated with cyclic-RGD to target cancer cells. The microspheres catalyze the formation of two anticancer agents 5FU and PP-121. The catalyzed reactions are propargyl cleavage and Suzuki-Miyaura cross-coupling. The dual-drug synthesis strategy was evaluated in brain cancer cells.

18. Miller MA, Askevold B, Mikula H, Kohler RH, Pirovich D, Weissleder R: **Nano-palladium is a cellular catalyst for in vivo chemistry.** *Nat Commun* 2017, **8**:15906.

••

Thorough study of a nanoencapsulated Pd-catalyst. The biocompatibility, localization in mice and use for drug release were investigated. The Pd-compound uncages doxorubicin and catalyzes a Heck coupling *in cellulo*.

19. Hrkach J, Von Hoff D, Mukkaram Ali M, Andrianova E, Auer J, Campbell T, De Witt D, Figa M, Figueiredo M, Horhota A, et al.: **Preclinical development and clinical translation of a PSMA-targeted docetaxel nanoparticle with a differentiated pharmacological profile.** *Sci Transl Med* 2012, **4**:128ra139.
20. Bai Y, Feng X, Xing H, Xu Y, Kim BK, Baig N, Zhou T, Gewirth AA, Lu Y, Oldfield E, et al.: **A Highly Efficient Single-Chain Metal-Organic Nanoparticle Catalyst for Alkyne-Azide "Click" Reactions in Water and in Cells.** *J Am Chem Soc* 2016, **138**:11077-11080.
21. Clavadetscher J, Hoffmann S, Lilienkamp A, Mackay L, Yusop RM, Rider SA, Mullins JJ, Bradley M: **Copper Catalysis in Living Systems and In Situ Drug Synthesis.** *Angew Chem Int Ed Engl* 2016, **55**:15662-15666.
22. Perez-Lopez AM, Rubio-Ruiz B, Sebastian V, Hamilton L, Adam C, Bray TL, Irusta S, Brennan PM, Lloyd-Jones GC, Sieger D, et al.: **Gold-Triggered Uncaging Chemistry in Living Systems.** *Angew Chem Int Ed Engl* 2017.

•

The Au-resin catalyzes the uncaging of several prodrugs in A549 cancer cell cultures. Next, the Au-resin was implanted into a zebrafish brain. The fluorophore rhodamine 110 was efficiently deprotected inside the brain.

23. Tonga GY, Jeong Y, Duncan B, Mizuhara T, Mout R, Das R, Kim ST, Yeh YC, Yan B, Hou S, et al.: **Supramolecular regulation of bioorthogonal catalysis in cells using nanoparticle-embedded transition metal catalysts.** *Nat Chem* 2015, **7**:597-603.

••

Pd- and Ru-complexes were encapsulated in Au-nanoparticles. The activity of these catalysts was controlled by a supramolecular 'gate-keeper'. Unmasking reactions were catalyzed after the addition

of the competitive activator. Rhodamine 110 and the cytotoxic drug 5FU were released in live HeLa cells. The reversible switch allowed a temporal controlled release of product.

24. Tornøe CW, Christensen C, Meldal M: **Peptidotriazoles on solid phase: [1,2,3]-triazoles by regioselective copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides.** *J Org Chem* 2002, **67**:3057-3064.
25. Rostovtsev VV, Green LG, Fokin VV, Sharpless KB: **A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective "ligation" of azides and terminal alkynes.** *Angew Chem Int Ed Engl* 2002, **41**:2596-2599.
26. Liang L, Astruc D: **The copper(I)-catalyzed alkyne-azide cycloaddition (CuAAC) "click" reaction and its applications. An overview.** *Coord Chem Rev* 2011, **255**:2933-2945.
27. Streu C, Meggers E: **Ruthenium-induced allylcarbamate cleavage in living cells.** *Angew Chem Int Ed Engl* 2006, **45**:5645-5648.
28. Volker T, Dempwolff F, Graumann PL, Meggers E: **Progress towards bioorthogonal catalysis with organometallic compounds.** *Angew Chem Int Ed Engl* 2014, **53**:10536-10540.
29. Volker T, Meggers E: **Chemical Activation in Blood Serum and Human Cell Culture: Improved Ruthenium Complex for Catalytic Uncaging of Alloc-Protected Amines.** *Chembiochem* 2017, **18**:1083-1086.
30. Sanchez MI, Penas C, Vazquez ME, Mascarenas JL: **Metal-catalyzed uncaging of DNA-binding agents in living cells.** *Chem Sci* 2014, **5**:1901-1907.
31. Hsu HT, Trantow BM, Waymouth RM, Wender PA: **Bioorthogonal Catalysis: A General Method To Evaluate Metal-Catalyzed Reactions in Real Time in Living Systems Using a Cellular Luciferase Reporter System.** *Bioconjug Chem* 2016, **27**:376-382.
32. Tomas-Gamasa M, Martinez-Calvo M, Couceiro JR, Mascarenas JL: **Transition metal catalysis in the mitochondria of living cells.** *Nat Commun* 2016, **7**:12538.
33. Murphy MP: **Targeting lipophilic cations to mitochondria.** *Biochim Biophys Acta* 2008, **1777**:1028-1031.
34. Smith RA, Porteous CM, Gane AM, Murphy MP: **Delivery of bioactive molecules to mitochondria in vivo.** *Proc Natl Acad Sci U S A* 2003, **100**:5407-5412.
35. Logan A, Pell VR, Shaffer KJ, Evans C, Stanley NJ, Robb EL, Prime TA, Chouchani ET, Cochemme HM, Fearnley IM, et al.: **Assessing the Mitochondrial Membrane Potential in Cells and In Vivo using Targeted Click Chemistry and Mass Spectrometry.** *Cell Metab* 2016, **23**:379-385.
36. Chalmers S, Caldwell ST, Quin C, Prime TA, James AM, Cairns AG, Murphy MP, McCarron JG, Hartley RC: **Selective uncoupling of individual mitochondria within a cell using a mitochondria-targeted photoactivated protonophore.** *J Am Chem Soc* 2012, **134**:758-761.
37. Sadhu KK, Eierhoff T, Romer W, Winssinger N: **Photoreductive uncaging of fluorophore in response to protein oligomers by templated reaction in vitro and in cellulo.** *J Am Chem Soc* 2012, **134**:20013-20016.
38. Sadhu KK, Lindberg E, Winssinger N: **In cellulo protein labelling with Ru-conjugate for luminescence imaging and bioorthogonal photocatalysis.** *Chem Commun (Camb)* 2015, **51**:16664-16666.
39. Holtzer L, Oleinich I, Anzola M, Lindberg E, Sadhu KK, Gonzalez-Gaitan M, Winssinger N: **Nucleic Acid Templated Chemical Reaction in a Live Vertebrate.** *ACS Cent Sci* 2016, **2**:394-400.

••

A Ru-complex was linked to PNA-probes targeting specific microRNAs. Rhodamine was released through photoreduction of an azide by the Ru-complex. The expression of miRNAs in specific tissues was monitored by fluorescence. This established a nucleic acid-templated chemical reaction in live vertebrates.

40. Jeschek M, Reuter R, Heinisch T, Trindler C, Klehr J, Panke S, Ward TR: **Directed evolution of artificial metalloenzymes for in vivo metathesis.** *Nature* 2016, **537**:661-665.

••

Binding of biotinylated Ru-complexes to streptavidin creates artificial metalloenzymes (ArMs). These ArMs catalyze an olefin metathesis in the periplasm of *E. coli*. The artificial metathase was improved through the directed evolution of streptavidin. The *in cellulo* catalysis facilitated an efficient and fast screening process. The evolved ArM outperforms the standard ring closing metathesis catalysts.

41. Li N, Lim RK, Edwardraja S, Lin Q: **Copper-free Sonogashira cross-coupling for functionalization of alkyne-encoded proteins in aqueous medium and in bacterial cells.** *J Am Chem Soc* 2011, **133**:15316-15319.
42. Liu CC, Schultz PG: **Adding new chemistries to the genetic code.** *Annu Rev Biochem* 2010, **79**:413-444.
43. Davis L, Chin JW: **Designer proteins: applications of genetic code expansion in cell biology.** *Nat Rev Mol Cell Biol* 2012, **13**:168-182.
44. Spicer CD, Triemer T, Davis BG: **Palladium-mediated cell-surface labeling.** *J Am Chem Soc* 2012, **134**:800-803.
45. Li J, Lin S, Wang J, Jia S, Yang M, Hao Z, Zhang X, Chen PR: **Ligand-free palladium-mediated site-specific protein labeling inside gram-negative bacterial pathogens.** *J Am Chem Soc* 2013, **135**:7330-7338.
46. Li N, Ramil CP, Lim RK, Lin Q: **A genetically encoded alkyne directs palladium-mediated protein labeling on live mammalian cell surface.** *ACS Chem Biol* 2015, **10**:379-384.
47. Li J, Yu J, Zhao J, Wang J, Zheng S, Lin S, Chen L, Yang M, Jia S, Zhang X, et al.: **Palladium-triggered deprotection chemistry for protein activation in living cells.** *Nat Chem* 2014, **6**:352-361.

••

The authors used Pd-complexes for gain-of-function studies in HeLa cells. Unnatural amino acids were encoded in OspF by amber stop codon suppression. The incorporation of a protected lysine in OspF inhibits its enzymatic activity. The enzyme was activated through cleavage of the protecting group. These gain of function approaches allow new functional studies.

48. Wang J, Zheng S, Liu Y, Zhang Z, Lin Z, Li J, Zhang G, Wang X, Li J, Chen PR: **Palladium-Triggered Chemical Rescue of Intracellular Proteins via Genetically Encoded Allene-Caged Tyrosine.** *J Am Chem Soc* 2016, **138**:15118-15121.
49. Indrigo E, Clavadetscher J, Chankeshwara SV, Megia-Fernandez A, Lilienkampf A, Bradley M: **Intracellular delivery of a catalytic organometallic complex.** *Chem Commun (Camb)* 2017, **53**:6712-6715.

•

A Pd-catalyst was conjugated to a cell-penetrating peptide. This new type of catalytic system has improved cell uptake rates. The Pd-complex unmasks rhodamine in prostate cancer cells.

50. Zahid M, Robbins PD: **Cell-type specific penetrating peptides: therapeutic promises and challenges.** *Molecules* 2015, **20**:13055-13070.
51. Svensen N, Walton JG, Bradley M: **Peptides for cell-selective drug delivery.** *Trends Pharmacol Sci* 2012, **33**:186-192.
52. Sasmal PK, Carregal-Romero S, Han AA, Streu CN, Lin Z, Namikawa K, Elliott SL, Koster RW, Parak WJ, Meggers E: **Catalytic azide reduction in biological environments.** *Chembiochem* 2012, **13**:1116-1120.
53. Bose S, Ngo AH, Do LH: **Intracellular Transfer Hydrogenation Mediated by Unprotected Organoiridium Catalysts.** *J Am Chem Soc* 2017, **139**:8792-8795.
54. Tsubokura K, Vong KK, Pradipta AR, Ogura A, Urano S, Tahara T, Nozaki S, Onoe H, Nakao Y, Sibgatullina R, et al.: **In Vivo Gold Complex Catalysis within Live Mice.** *Angew Chem Int Ed Engl* 2017, **56**:3579-3584.

••

Au-complexes were exploited to specifically label intestine and liver of live mice. The Au-complex was targeted to these organs by linking it to coumarin. This coumarin-Au-complex binds with high affinity to albumins. In turn, the albumins were directed to liver and intestine by an N-glycan coating.